



Measure Specific Guidelines for Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD)

Description:

The percentage of members 18–64 years of age with schizophrenia, schizoaffective disorder or bipolar disorder, who were dispensed an antipsychotic medication and had a diabetes screening test during the measurement year.

Diabetes Screening is defined as:

A glucose test or HbA1c test performed during the measurement year

Measure Requirements:

- Members with any diagnosis of schizophrenia, schizoaffective disorder, or bipolar disorder during the measurement year*
- AND**
- Evidence of diabetes screening (glucose test or HbA1c test) performed during the measurement year

**Diagnosis is obtained through claim/encounter data*

Exclusions for the measure:

- Members receiving hospice services
- Members with a diagnosis of diabetes*
- Members who were dispensed insulin or oral hypoglycemics/antihyperglycemics during the measurement year*
- Members who had no antipsychotic medication dispensed during the measurement year*

**Diagnosis and dispensing events obtained through claim/encounter or pharmacy data*

Tips for compliance:

- ✓ Documentation of the type of diabetes screening and the date it was completed
- ✓ The use of Current Procedural Terminology (CPT) and/or Category II codes for evidence of services may decrease the need for record abstraction and chart review, thereby minimizing administrative burden:

Diabetes Screening	
CPT	CPT Definition
80047 80048 80050 80053 80069 82947 82950 82951	Glucose Lab Test
83036 83037	HbA1c Lab Test
CPT-CAT-II	CPT-CAT-II Definition
3044F	Most recent hemoglobin A1c (HbA1c) level less than 7.0% (DM)
3046F	Most recent hemoglobin A1c level greater than 9.0% (DM)
3051F	Most recent hemoglobin A1c (HbA1c) level greater than or equal to 7.0% and less than 8.0% (DM)
3052F	Most recent hemoglobin A1c (HbA1c) level greater than or equal to 8.0% and less than or equal to 9.0% (DM)